

APORPHINE ALKALOIDS FROM *Ocotea macrophylla* (LAURACEAE)

Ludy Cristina Pabon\* y Luis Enrique Cuca

Departamento de Química, Facultad de Ciencias, Universidad Nacional de Bogotá, KR 30 45 03, Colombia. AA 14490

Recebido em 26/6/09; aceite em 6/11/09; publicado na web em 10/3/10

Four aporphine alkaloids from the wood of *Ocotea macrophylla* (Lauraceae) were isolated and characterized as (*S*)-3-methoxy-nordomesticine (**1**), (*S*)-*N*-ethoxycarbonyl-3-methoxy-nordomesticine (**2**), (*S*)-*N*-formyl-3-methoxy-nordomesticine (**3**) and (*S*)-*N*-methoxycarbonyl-3-methoxy-nordomesticine (**4**); alkaloids 2-4 are being report for the first time. The structure the isolated compounds were determined based on their spectral data and by comparison of their spectral data with values described in literature. The alkaloid fraction and compound **1** showed antifungal activity against *Fusarium oxysporum* f. sp. *lycopersici* and also compound **1** showed antimicrobial activity towards *Staphylococcus aureus*, *Enterococcus faecalis* as well.

Keywords: *Ocotea macrophylla*; aporphine alkaloids; Lauraceae.

## INTRODUCTION

The *Ocotea* genus (Lauraceae) includes more than 350 species found in the American continent and southern Africa.<sup>1,2</sup> In Colombia, there are 35 *Ocotea* species distributed throughout the country mainly in the Andean forests.<sup>3</sup> In traditional medicine, some *Ocotea* species shown different applications. *O. quixos* is used as disinfectant, local anesthetic and anti-diarrheic.<sup>4</sup> *O. lancifolia* is used as antiparasitic, and *O. caparrapi* is used to treat insect bites, snake bites, bronchitis, and cancerous tumors.<sup>5,6</sup>

Chemically, the *Ocotea* genus is known mainly as a source of metabolites type furofuran<sup>7</sup> and tetrahydrofuran lignans,<sup>8</sup> bicyclo[3.2.1] octane<sup>9</sup> and benzofuran neolignans,<sup>10</sup> and benzyloquinoline<sup>11</sup> and aporphine alkaloids.<sup>5</sup> In previous studies, four aporphine alkaloids from wood of *Ocotea macrophylla* were isolated and identified as nantenine, glaucine, isocorydine and dehydronantenine.<sup>12,13</sup> In this paper, we describe isolation and structural determination of three new aporphine alkaloids **2-4** besides (*S*)-3-methoxy-nordomesticine **1** and the reports antibacterial and antifungal activities of compounds.

## RESULT AND DISCUSSION

The ethanolic extract from the stem of *O. macrophylla* was subjected to an acid-base extraction to obtain an alkaloidal fraction, that was further subjected to fractionation and purification by chromatographic methods leading to the isolation of four alkaloids: (*S*)-3-methoxy-nordomesticine (**1**), (*S*)-*N*-ethoxycarbonyl-3-methoxy-nordomesticine (**2**), (*S*)-*N*-formyl-3-methoxy-nordomesticine (**3**) and (*S*)-*N*-methoxycarbonyl-3-methoxy-nordomesticine (**4**). The structures of the alkaloids **1-4** are shown in Figure 1 and spectroscopic data in the Table 1.

Alkaloid **1** was obtained as a yellowish oil that gives a positive reaction to Dragendorff reagent, which suggests the presence of an alkaloid. The molecular formula of **1** was established as C<sub>19</sub>H<sub>19</sub>NO<sub>5</sub> from HRESIMS at *m/z* 340.1178 [M-H]<sup>-</sup>, indicating eleven degrees of insaturation. Its UV spectrum showed maximum absorptions at 224 and 308 nm, characteristic of aporphine alkaloids,<sup>14</sup> and the IR spectrum displayed hydroxyl (3404 cm<sup>-1</sup>), aromatic (1585 and 1467 cm<sup>-1</sup>), and methylenedioxy (1039 and 935 cm<sup>-1</sup>) absorptions.<sup>15</sup> The <sup>1</sup>H NMR spectrum shows typical signals for aporphine alkaloids at:




Figure 1. Basic structure of isolated aporphine alkaloids

$\delta$  4.74 (1H, *dd*,  $J=3.9$  and 13.8 Hz, H-6a), 3.73 (1H, *ddd*,  $J=12.1$ , 4.3 and 2.1 Hz, H-5), 3.24 (1H, *ddd*,  $J=12.1$ , 8.4 and 3.6 Hz, H-5), 2.93 (2H, *m*, H-3, H-7) and 2.64 (2H, *m*, H-3, H-7). Also, two singlets were present in the aromatic region at  $\delta$  7.90 (H-11), 6.70 (H-8) besides the hydroxyl group at 6.30, as well as signals for a methylenedioxy group at  $\delta$  5.95 (1H, *d*,  $J=1.3$  Hz) and 5.94 (1H, *d*,  $J=1.3$  Hz)<sup>16</sup> and two methoxyl groups at  $\delta$  3.97 and 3.87 (3H, each *s*). The <sup>13</sup>C NMR and DEPT showed spectra of **1** showed 19 signals, corresponding to two methoxyl groups at  $\delta$  60.9 and 60.3; four methylene groups at  $\delta$  100.9, 43.9, 35.1 and 23.8, where the first signal corresponds to a methylenedioxy group; three methines at  $\delta$  108.5, 108.6, and 52.8, and ten quaternary carbons, whose chemical shift are found in Table 1.

The signal at  $\delta_{\text{H}}$  6.30 (1H, *s*) showed no connectivity in the HMQC experiment, indicative of the presence of a phenolic OH group, supported by IR. Analysis of the HMBC spectrum allowed the location of substituents, according to the correlations observed between the signals at  $\delta_{\text{H}}$  5.94 and 5.95 (for the methylenedioxy group) with the signals at  $\delta_{\text{C}}$  145.9 (C-9) and 146.2 (C-10), and these last two signals with the protons at  $\delta_{\text{H}}$  6.70 (H-8) and 7.90 (H-11), respectively, suggesting the presence of a methylenedioxy group at positions 9 and 10 and two hydrogens on the aromatic ring in a *para* orientation. The presence of the hydroxyl group at position 1, was determined using correlations between the signals at  $\delta_{\text{H}}$  6.30 and  $\delta_{\text{C}}$  116.5, assigned to C-11b. The location of the methoxyl groups in positions C-2 and C-3, were assigned according to the correlation of

\*e-mail: lcpabonb@unal.edu.co

**Table 1.**  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopic assignments of **1-4** in ( $\text{CDCl}_3$ ,  $^1\text{H}$  NMR: 400 MHz and  $^{13}\text{C}$  NMR: 100 MHz)

Position	Alkaloid 1		Alkaloid 2		Alkaloid 3a		Alkaloid 3b		Alkaloid 4	
	$\delta_{\text{H}}$ (mult., J, Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (mult., J, Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (mult., J, Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (mult., J, Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (mult., J, Hz)	$\delta_{\text{C}}$
1		144.8		144.8		145.2		145.0		144.8
2		138.4		138.5		139.0		138.7		138.5
3		147.8		147.9		147.8		148.3		147.9
3a		118.5		119.4		118.3		119.3		119.3
4	2.93 (m) 2.64 (m)	23.8	2.90 (m) 2.53 (m)	23.5	3.04 (m)	24.1	3.04 (m)	23.0	2.95 (m) 2.52 (m)	23.4
5	3.73 (ddd, 12.1, 4.3, 2.1) 3.24 (ddd, 12.1, 8.4, 3.6)	43.9	4.45 (d, 11.8) 2.95 (m)	38.6	3.82 (ddd, 12.7, 4.7, 1.7) 3.31 (dt, 12.7, 2.7)	41.8	3.05 (m) 2.72 (m)	37.0	4.43 (m) 2.87 (m)	38.7
6a	4.74 (dd, 13.8, 3.9)	52.8	4.76 (dd, 13.5, 4.2)	51.8	4.94 (dd, 13.6, 4.2)	49.6	4.46 (m)	53.7	4.71 (d, 11.2)	51.8
7	2.93 (m) 2.64 (m)	35.1	2.78 (m)	34.8	2.95 (m) 2.78 (m)	33.7	3.04 2.66 (m)	35.9	2.95 (m) 2.52 (m)	34.0
7a		130.0		130.2		129.6		128.8		130.1
8	6.70 (s)	108.5	6.74 (s)	108.5	6.76 (s)	108.8	6.74 (s)	108.4	6.74 (s)	108.5
9		145.9		146.0		146.1		145.2		145.9
10		146.2		146.3		146.4		147.7		146.2
11	7.90 (s)	108.6	7.94 (s)	108.8	7.93 (s)	108.7	7.94 (s)	108.9	7.94 (s)	108.7
11a		125.2		125.3		125.0				125.3
11b		116.5		116.4		116.6		117.3		116.3
11c		129.9		129.1		128.8				128.9
OH	6.30 (s)		6.32 (s)		6.33 (s)		6.33 (s)		6.44 (s)	
2-OCH <sub>3</sub>	3.97 (s)	60.9	3.96 (s)	61.0	3.96 (s)	61.0	3.97 (s)	61.0	3.95 (s)	60.9
3-OCH <sub>3</sub>	3.87 (s)	60.3	3.86 (s)	60.4	3.86 (s)	60.5	3.86 (s)	60.4	3.85 (s)	60.4
O-CH <sub>2</sub> -O	5.95 (d, 1.3) 5.94 (d, 1.3)	100.8	5.97 (d, 1.4) 5.96 (d, 1.4)	100.9	5.97 (d, 1.3) 5.96 (d, 1.3)	100.9	5.99 (d, 1.4) 5.97 (d, 1.4)	101.0	5.96 (d, 1.3) 5.95 (d, 1.3)	100.8
C=O				158.8	8.25 (s)	161.9	8.37 (s)	161.8		155.8
-OCH <sub>3</sub>			4.23 (m)	61.0						
-OCH <sub>2</sub>									3.76 (s)	52.6
CH <sub>3</sub>			1.29 (t, 7.10)	14.8						

*J* values (Hz) are in parenthesis.

the hydrogens at  $\delta_{\text{H}}$  3.96 and 3.86 with the carbons at  $\delta_{\text{C}}$  138.4 (C-2) and  $\delta_{\text{C}}$  147.8 (C-3), respectively.

The absolute configuration of C-6a was assigned as *S*, because it has a negative Cotton effect at 280 nm and a positive Cotton effect at 240 nm in CD curve.<sup>17</sup> Additionally, this was confirmed by the positive value of optical rotation  $[\alpha]_{\text{D}}^{25} = +51.7$  (*c* 0.38,  $\text{CHCl}_3$ ).<sup>18</sup> Therefore, alkaloid **1** was determined as (*S*)-3-methoxy-nordomesticine, an aporphine alkaloid reported previously from *Nectandra sinuata*<sup>19</sup> also belonging to the Lauraceae family. This report corrects and completes spectroscopic data for this compound.

Alkaloid **2** was obtained as a yellow oil that gives a positive reaction to Dragendorff reagent and its optical rotation value was  $[\alpha]_{\text{D}}^{25} [\alpha]_{\text{D}}^{25} = +33.3$  (*c* 0.60,  $\text{CHCl}_3$ ). The UV and IR spectrum of **2** were similar to those of **1**, except for the appearance, in both spectra, of absorptions due to a carbamate group at 282 nm and at 1688  $\text{cm}^{-1}$ , respectively.<sup>20</sup> The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra showed a similar profile to **1**. In the  $^1\text{H}$  NMR two new signals appeared at  $\delta_{\text{H}}$  4.29 (2H, *m*) and 1.29 (3H, *m*), as well as the displacement of the signal H-5, due to the presence of a deprotecting group in proximity. The COSY experiment showed the correlation of the signals  $\delta_{\text{H}}$  4.29 and 1.29, which indicates the presence of an ethyl group that due to its displacement, suggested to be attached to a heteroatom. The  $^{13}\text{C}$  NMR and DEPT

spectra showed the appearance of three new signals at  $\delta_{\text{C}}$  158.8 (C), 61.0 ( $\text{CH}_2$ ), and 14.8 ( $\text{CH}_3$ ), typical signals of an ethoxycarbonyl group attached to a nitrogen atom. The absolute configuration of C-6a was determined as *S*, because it showed the same Cotton effects as **1**. Therefore the alkaloid compound **2** was identified as (*S*)-*N*-ethoxycarbonyl-3-methoxy-nordomesticine. Its ESIMS spectrum gave a pseudomolecular ion peak at  $m/z$  414  $[\text{M}+\text{H}]^+$  corresponding to the molecular formula of  $\text{C}_{22}\text{H}_{22}\text{NO}_7$ , and fragmentations were due to the loss of  $\text{CH}_3\text{OH}$  and  $\text{CH}_3\text{CH}_2\text{OH}$  at  $m/z$  382  $[\text{M}+\text{H}-32]^+$  and  $m/z$  368  $[\text{M}+\text{H}-46]^+$ , respectively. The negative ion mode ESI spectrum showed peaks at  $m/z$  412  $[\text{M}-\text{H}]^-$  and  $m/z$  383  $[\text{M}-\text{H}-29]^-$ , being the last one the loss of an ethyl group. This type of alkaloids with substituents *N*-ethoxycarbonyl have been isolated from *Lindera angustifolia* (Lauraceae).<sup>21</sup>

Alkaloid **3**, was obtained as a yellow oil, with an optical rotation value of  $[\alpha]_{\text{D}}^{25} [\alpha]_{\text{D}}^{25} = +7.5$  (*c* 0.53  $\text{CHCl}_3$ ). The IR spectrum was similar to that of alkaloids **1** and **2**. Additionally, there were observed the presence of absorptions at 2830, 2700, 1739  $\text{cm}^{-1}$ , characteristic for formyl group. In HRESIMS negative mode was observed the pseudomolecular ion  $[\text{M}-\text{H}]^-$  at  $m/z$  369.1133, corresponding to the molecular formula of  $\text{C}_{20}\text{H}_{20}\text{NO}_6$ . The negative ion mode of ESI-MS spectra showed the loss of a formyl group at  $m/z$  339  $[\text{M}-\text{H}-29]^-$ . The



2923, 2855, 1585, 1464, 1274, 1129, 1041, 935, 757  $\text{cm}^{-1}$ ; CD  $[\theta]_{242} +3345$ ,  $[\theta]_{280} -250$ ; for  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy data, see Table 1; negative ESIMS:  $m/z = 340$   $[\text{M}-\text{H}]^-$ ; HR-ESIMS:  $m/z = 340.1178$   $[\text{M}-\text{H}]^-$  (calcd. 340.1185 for  $\text{C}_{19}\text{H}_{19}\text{NO}_3$ ).

(*S*)-*N*-Ethoxycarbonyl-3-methoxy-nordomesticine (**2**): Yellow oil;  $[\alpha]_D^{25} = +33.3$  ( $c$  0.60,  $\text{CHCl}_3$ ); UV (MeOH) 220, 282, 308 nm; IR (film):  $\nu_{\text{max}}$  3402, 2928, 2830, 2700, 1739, 1464, 1236, 1145, 1039, 936, 757  $\text{cm}^{-1}$ ; CD  $[\theta]_{238} +3634$ ,  $[\theta]_{280} -100$ ; for  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy data, see Table 1; positive ESIMS:  $m/z = 414$   $[\text{M}+\text{H}]^+$ , 382  $[\text{M}+\text{H}-\text{CH}_3\text{OH}]^+$ , 368  $[\text{M}+\text{H}-\text{CH}_2\text{CH}_3\text{OH}]^+$ ; negative ESIMS:  $m/z = 412$   $[\text{M}-\text{H}]^-$ , 383  $[\text{M}-\text{H}-\text{CH}_2\text{CH}_3]^-$ ; HR-ESIMS:  $m/z = 412.1385$   $[\text{M}-\text{H}]^-$  (calcd. 412.1396 for  $\text{C}_{22}\text{H}_{22}\text{NO}_7$ ).

(*S*)-*N*-Formyl-3-methoxy-nordomesticine (**3**): Yellow oil;  $[\alpha]_D^{25} = +7.5$  ( $c$  0.53, DMSO); UV (MeOH) 220, 282, 310 nm; IR (film):  $\nu_{\text{max}}$  3403, 2925, 1688, 1464, 1270, 1145, 1039, 936, 768  $\text{cm}^{-1}$ ; CD  $[\theta]_{242} +2321$ ,  $[\theta]_{280} -250$ ; for  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy data, see Table 1; positive ESIMS:  $m/z = 369$   $[\text{M}+\text{H}]^+$ ; negative ESIMS:  $m/z = 368$   $[\text{M}-\text{H}]^-$ , 339  $[\text{M}-\text{H}-\text{CHO}]^-$ ; HR-ESIMS:  $m/z = 369.1133$   $[\text{M}-\text{H}]^-$  (calcd. 369.1134 for  $\text{C}_{20}\text{H}_{18}\text{NO}_6$ ).

(*S*)-*N*-Methoxycarbonyl-3-methoxy-nordomesticine (**4**): Solid cream, melting point 222-223°C (MeOH);  $[\alpha]_D^{25} = +47.0$  ( $c$  0.55,  $\text{CHCl}_3$ ); UV (MeOH) 220, 282, 308 nm; IR (film):  $\nu_{\text{max}}$  3372, 2927, 1685, 1466, 1272, 1146, 1039, 935, 876  $\text{cm}^{-1}$ ; CD  $[\theta]_{239} +303$   $[\theta]_{280} -10$ ; for  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy data, see Table 1; positive ESIMS:  $m/z = 400$   $[\text{M}+\text{H}]^+$ , 368  $[\text{M}+\text{H}-\text{CH}_3\text{OH}]^+$ ; negative ESIMS:  $m/z = 398$   $[\text{M}-\text{H}]^-$ , 339  $[\text{M}-\text{H}-\text{CH}_3\text{OCO}]^-$ ; HR-ESIMS:  $m/z = 398.1233$   $[\text{M}-\text{H}]^-$  (calcd. 398.1240 for  $\text{C}_{21}\text{H}_{20}\text{NO}_7$ ).

### Antifungal assay

*F. oxysporum* was obtained from the culture collection of University of Cundinamarca (Department of Agronomy, Laboratory of Phytopathology). PDA was used as medium for antifungal activity assays. The culture medium was inoculated with 100  $\mu\text{L}$  of a solution of  $10^5$  spores. The samples were prepared in solutions of different concentrations, corresponding to 50, 25, and 10  $\mu\text{g}/\mu\text{L}$  of the alkaloid fraction and 5, 2.5, 1.0, 0.5, 0.2, and 0.1  $\mu\text{g}/\mu\text{L}$  of the pure alkaloids. Ten microliters of samples were applied on the filter paper discs and placed on the inoculated medium. The plates were sealed and left in an incubator for 3 days at 25 °C. Clear zones appearing against a growing fungus indicated the minimal amount of fraction or alkaloid required to inhibit the fungi growth. Three replicates were made for each treatment. Benomyl (benzimidazole – 5  $\mu\text{g}$ ) was used as a positive control, and acetone served as a negative control.<sup>23</sup>

### Antibacterial assays

The antibacterial activity was evaluated by radial diffusion method adapted from the methodology previously published by Lehrer.<sup>27</sup> The compounds were evaluated against two Gram (+) strains: *Staphylococcus aureus* ATCC 6538 and *Streptococcus fecalis* ATCC 29212 and three Gram (-) strains: *Escherichia coli* ATCC 25922 and *Salmonella tiphymurium*, ATCC 14028s and *Salmonella tiphymurium* MS7953. A colony isolated from each strain, was deposited in 3 mL of soy trypticase (TSB) for Gram (+) strains and Luria Broth (LB) for Gram (-) strains, and were incubated at 37 °C with stirring, until the microorganisms were in the logarithmic phase. The supernatant was removed and the sediment obtained was re-suspended in phosphate buffer (PBS), followed by washes with PBS and centrifuging. Finally the sediment was re-suspended

in PBS and the optical density determined in 620 nm to calculate the number of CFU (colony forming units) per milliliter. It disperses a certain volume that contains  $4 \times 10^7$  CFU in each dish. The measured volume was mixed and homogenized in 15 mL agarose fused to more or less 45 °C. This bacterial suspension was served in petri dishes and left to solidify at room temperature, after which were made of 2 mm diameter holes with a sterile punch.

The test samples were prepared dissolving 1 mg of the pure compound in 500  $\mu\text{L}$  of DMSO, which are placed 8  $\mu\text{L}$  of the sample in duplicate and incubated at 37 °C for 30 min. After this time the nutrient medium was added, which contains molten agar agar and TSB, incubated for 18 h at 37 °C and then diameter of inhibition zones was measured by the activity of the compound. Positive controls used were different antibiotics, Ampicillin (50 mg/mL), Kanamycin (10 mg/mL) and Tetracycline (4.12 mg/mL) at a dilution 1:100 in PBS and was used as negative controls DMSO and PBS, each control 8  $\mu\text{L}$  be served by each well. The diameters of inhibition zones were measured in millimeters and the results were reported as units of activity according to the ratio which stipulates that 1 Unit of Action (UA) is equal to 0.1 mm of the inhibition zone.

### SUPPLEMENTARY MATERIAL

1D and 2D NMR spectra for compounds **1-4**. This material is available free of charge at <http://quimicanova.sbq.org.br>, in .PDF format.

### ACKNOWLEDGMENTS

The authors would like to thank the National University of Colombia for its financial support for this project. Also, the Nuclear Magnetic Resonance Laboratory and the mass spectrometry laboratory are acknowledged for their support in recording the NMR spectra and the HR-ESI-MS, respectively. Furthermore, the authors would like to thank the Javeriana University for the optical rotations, the FIDIC (Foundation Institute of Immunology of Colombia) for recording the CD spectra and especially to Prof. J. M. Lozano by assays antibacterial activity.

### REFERENCES

- Takaku, S.; Haber, W.; Setzer, W.; *Biochem. Syst. Ecol.* **2007**, *35*, 525.
- Guerrini, A.; Sacchetti, G.; Muzzolli, M.; Moreno, G.; Medici, A.; Besco, E.; Bruni R.; *J. Agric. Food Chem.* **2006**, *54*, 7778.
- IDEAM Distribución de 15 familias forestales en Colombia. Atlas interactivo, <http://www.ideam.gov.co/atlas/mecosis.htm>, accessed in March 2008.
- Ballabeni, V.; Tognoli, M.; Bertoni, S.; Bruni, R.; Guerrini, A.; Moreno, G.; Barocelli, E.; *Pharmacol. Res.* **2007**, *55*, 23.
- Fournet, A.; Ferreira, M.; Rojas, A.; Guy, I.; Guinaudeau, H.; Heinzen, H.; *Fitoterapia.* **2007**, *78*, 382.
- Palomino, E.; Maldonado, C.; *J. Nat. Prod.* **1996**, *59*, 77.
- Marques, R.; Batistuzzo, S.; Silva, C.; Barbosa, J.; Fassarella, L.; *Mutat. Res.* **2003**, *536*, 117.
- López, H.; Valera, A.; *J. Nat. Prod.* **1995**, *58*, 782.
- Zschocke, S.; Staden, J.; Paulus, K.; Bauer, R.; Horn, M.; Munro, O.; Brown N.; Drewes, S.; *Phytochemistry* **2000**, *54*, 591.
- Lordello, A.; Yoshida, M.; *Phytochemistry* **1997**, *46*, 741.
- Silva, I.; Barbosa, J.; Silva, M.; Lacerda C. da-Cunha, E.; *Biochem. Syst. Ecol.* **2002**, *30*, 881.
- Franca, N.; Giesbrecht, A.; Gottlieb, O.; Malgalhaes, A.; Malgalhaes, E.; Maia, J.; *Phytochemistry* **1975**, *14*, 1671.
- Warrumby, S.; Lordello, A.; *Quim. Nova* **2007**, *30*, 92.
- Wu, Y.; Chao, Y.; Chang, F.; Chen, Y.; *Phytochemistry* **1996**, *46*, 181.

15. Mathouet, H.; Elomri, A.; Lameiras, P.; Daich, A.; Vérité, P.; *Phytochemistry* **2007**, *68*, 1813.
16. Tantisewie, B.; Pharadai, T.; Pandhuganont, M.; Guinaudeau, H.; Freyer, A.; Shamma, M.; *J. Nat. Prod.* **1989**, *52*, 652.
17. Ringdahl, B.; Chan, R.; Craig, J.; *J. Nat. Prod.* **1981**, *44*, 80.
18. Nishiyama, Y.; Moriyasu, M.; Ichimaru, M.; Iwasa, K.; Kato, A.; Mathenge, S.; Chalo, P.; Juma, F.; *Phytochemistry* **2006**, *67*, 2671.
19. Castro, O.; Hasbun, C.; Calderon, M.; *Fitoterapia* **1991**, *62*, 72.
20. Chen, Y.; Chang, F.; Wu, Y.; *J. Nat. Prod.* **1996**, *59*, 904.
21. Zhao, O.; Zhao, Y.; Wang, K.; *J. Ethnopharm.* **2006**, *106*, 408.
22. Chang, F.; Wei, J.; Teng, C.; Wu, Y.; *J. Nat. Prod.* **1998**, *61*, 1457.
23. Gomez, Y.; Gil, K.; González, E.; Farías, L.; *Rev. Biol. Trop.* **2007**, *55*, 767.
24. Kuo, R.; Chang, F.; Chen, C.; Teng, C.; Yen, H.; Wu, Y.; *Phytochemistry* **2001**, *57*, 421.
25. Agnihotri, V.; ElSohly, H.; Khan, S.; Jacob, M.; Joshi, V.; Smillie, T.; Khan, I.; Walker, L.; *Phytochem. Lett.* **2008**, *1*, 89.
26. Ma, W.; Fukuski, Y.; Tahara, S.; Osawa, T.; *Fitoterapia* **2000**, *71*, 527.
27. Lehrner R.; Rosenman, M.; Harwig, S.; Jackson, R.; Eisenhaber, P.; *J. Immunol. Methods.* **1991**, *137*, 167.
28. Nimgirawath, S.; Udomputtimekakul, P.; Taechowisan, T.; Wanbanjob, A.; Shen Y.; *Chem. Pharm. Bull.* **2009**, *57*,