

CALDEN SINIC ACID, A BENZOIC ACID DERIVATIVE AND OTHERS COMPOUNDS FROM *Piper carniconnectivum*

Harley da Silva Alves, Maria de Fátima Vanderlei de Souza e Maria Célia de Oliveira Chaves*

Laboratório de Tecnologia Farmacêutica, Universidade Federal da Paraíba, CP 5009, 58051-970 João Pessoa - PB, Brasil




Figure 1S. IR spectrum (λ_{max} , KBr, cm^{-1}) of 1

*e-mail: cchaves@lft.ufpb.br




Figure 2S. ^1H NMR spectrum of **1** (δ , 500 MHz, CD_3CN)




Figure 3S. Expansion of the ^1H NMR spectrum of 1 (δ , 500 MHz, $\text{C}_3\text{D}_5\text{N}$)




Figure 4S. ^{13}C NMR spectrum of **1** (δ , 125 MHz, $\text{C}_3\text{D}_3\text{N}$)




Figure 5S. Expansion of the ^{13}C NMR of 1 (δ , 125 MHz, $\text{C}_5\text{D}_5\text{N}$)




Figure 6S. Expansion of the $^1\text{H} \times ^{13}\text{C}$ -HSQC NMR spectrum of **1**




Figure 7S. $^1\text{H} \times ^{13}\text{C}$ -HMBC NMR spectrum of 1




Figure 8S. Expansion of the $^1\text{H} \times ^{13}\text{C}$ - HMBC NMR spectrum of **1**




Figure 9S. Expansion of the $^1\text{H} \times ^{13}\text{C}$ - HMBC NMR spectrum of **1**




Figure 10S. $^1\text{H} \times ^1\text{H}$ - COSY NMR spectrum of **1**




Figure 11S. $^1\text{H} \times ^1\text{H}$ – NOESY NMR spectrum of **1**




Figure 12S. MS spectrum of **1**