

TRITERPENES AND NEW SAPONINS FROM *Ilex chamaedryfolia*: CHEMOTAXONOMIC TOOL TO *Ilex* SPECIES DIFFERENTIATION

Claiton L. Lencina, Mariana C. de Cardoso, Ivomar Zancanaro e Grace Gosmann*

Faculdade de Farmácia, Universidade Federal do Rio Grande do Sul, Av. Ipiranga, 2752, 90610-000 Porto Alegre - RS, Brasil

Viviane S. Pires, Pascal Sonnet e Dominique Guillaume#

UMR-CNRS 6219, Laboratoire des glucides, Faculté de Pharmacie, Université de Picardie Jules Verne, 1 rue des Louvels, 80037 Amiens, France

Eloir P. Schenkel

Centro de Ciências da Saúde, Universidade Federal de Santa Catarina, 88040-900 Florianópolis - SC, Brasil

Recebido em 16/2/10; aceito em 13/9/10; publicado na web em 8/12/10

Three saponins were isolated from leaves of *Ilex chamaedryfolia*. Their structures were established by spectroscopic and mass spectrometry data as the new saponin 3 β -*O*- β -D-glucopyranosyl-(1-3)- α -L-arabinopyranosyl-20(*S*)-19 α -hydroxyurs-12-en-28-oic acid 28-*O*- β -D-glucopyranosyl-(1-3)- β -D-glucopyranosyl ester, the new saponin 3 β -*O*- β -D-glucopyranosyl-(1-3)- α -L-arabinopyranosyl-20(*S*)-19 α -hydroxyurs-12-en-28-oic acid 28-*O*- β -D-glucopyranosyl ester and the known saponin 3 β -*O*- β -D-glucuronopyranosyl-20(*R*)-19 α -hydroxyurs-12-en-28-oic acid 28-*O*- β -D-glucopyranosyl ester. Ursolic acid and α -amyryn were also isolated.

Keywords: *Ilex chamaedryfolia*; Aquifoliaceae; triterpenes.

INTRODUCTION

We initiated some years ago a program aimed to identified saponins from South American *Ilex* species¹ considering the importance of this genera which includes *Ilex paraguariensis* A. St. Hil., named yerba mate. This species is widely used in South Brazil, Argentina, Paraguay and Uruguay to obtain the raw material used to prepare the traditional beverage called maté which is a very important economical crop. Hence, we already reported the structure of several saponins isolated from maté and other South American *Ilex* species: *I. affinis*, *I. argentina*, *I. brevicuspis*, *I. buxifolia*, *I. dumosa*, *I. integerrima*, *I. microdonta*, *I. psammophila*, *I. pseudobuxus*, *I. taubertiana* and *I. theezans*.¹⁻⁸

Ilex chamaedryfolia Reissek is a native shrub in Southern Brazil, found in the states of Rio Grande do Sul, Paraná and Santa Catarina, where it is known as “congonha miúda”, “congonha” or “congonha brava”.^{2,9} *I. chamaedryfolia* is one of the species eventually reported as an adulterant of the *I. paraguariensis*. Herein it is described the isolation and structural elucidation of three saponins and two triterpenes (ursolic acid and α -amyryn) from the leaves of *I. chamaedryfolia*. As far as we know, the saponins reported herein have not been yet described in the literature.

EXPERIMENTAL

Plant material

Leaves from *I. chamaedryfolia* Reissek were collected in Guaratuba, State of Paraná, Brazil. A specimen is on deposit in the Herbarium of the Department of Botany (ICN) at the Universidade

Federal do Rio Grande do Sul, Porto Alegre, Brazil (M. Sobral and E. P. Santos 9308).

General

Optical Rotation was measured on a Perkin-Elmer[®] 341 polarimeter. FAB-MS analysis was performed in positive mode on a Kratos MS 80 instrument and HRMS MS spectra were recorded on a Q-ToF micro Waters[®] high resolution mass spectrometer, operating on electrospray ionization mode. NMR spectra (¹H, 500 MHz; ¹³C, 125 MHz) were recorded in CD₃OD and CDCl₃ on a Bruker[®] Avance 500 spectrometer. Thin-layer chromatography (TLC) was on Si gel GF₂₅₄ Merck or Aldrich[®] using CH₂Cl₂:MeOH (98:2, v/v) and CHCl₃:MeOH:H₂O (80:40:5, v/v) as eluents for sapogenins and saponins, respectively. Compounds were visualised using anisaldehyde-sulphuric acid reagent and heating (100 °C).

Extraction and isolation

Air-dried powdered leaves (475 g) were submitted to maceration in aqueous 80% EtOH (1:10 plant:solvent, m/v) at room temperature (2 x 10 days). The ethanol extract was evaporated to dryness under reduced pressure and the residue (85 g, 18%) was suspended in water (1500 mL) and extracted successively with dichloromethane (3 x 500 mL), ethyl acetate (3 x 500 mL) and *n*-butanol (3 x 500 mL). Each organic phase was separately evaporated to obtain 5 g (0.7%), 5 g (0.7%) and 45 g (9.5%) of each fraction, respectively. The aqueous residue was lyophilised to obtain 27 g (5.6%).

Dichloromethane fraction (1 g) was submitted to CC on silica gel eluting with gradients of cyclohexane:ethyl acetate mixtures. Fractions were pooled according to TLC. Fractions 35-41 (88 mg) and 82-85 (30 mg) were subjected to crystallisation in yielding compounds **1** (19 mg) and **2** (25 mg) in a pure form, respectively. Part of the *n*-butanol fraction (10 g) was submitted to successive CC on silica

*e-mail: grace.gosmann@ufrgs.br

#Present address: Faculté de Pharmacie, Université de Reims, Chimie Thérapeutique, 51 rue Cognacq Jay, 51100 Reims, France

gel eluting with gradients of CH₂Cl₂:EtOH:H₂O mixtures, or CC on LiChroprep™ RP-18 using H₂O:MeOH. Fractions were pooled according to their TLC profiles. Compounds **3** (13 mg), **4** (10 mg) and **5** (13 mg) were isolated (Figure 1). One fraction containing mainly compound **3** was acetylated⁶ to obtain pure **3a** (14 mg).

α-amyrin (1). This compound was identified after comparison of its physical-chemical data with the literature¹⁰ and by co-TLC with an authentic sample.

Ursolic acid (2). (3β)-3-hydroxyurs-12-en-28-oic acid. It was identified by co-TLC with an authentic sample.¹¹

Compound 3 was identified as the deacylated form of **3a** after comparison of both NMR data.

Peracetylated 3β-O-β-D-glucopyranosyl-(1-3)-α-L-arabinopyranosyl-20(S)-19α-hydroxyurs-12-en-28-oic acid 28-O-β-D-glucopyranosyl-(1-3)-β-D-glucopyranosyl ester (3a). Amorphous powder. FABMS (positive mode) *m/z*: 1659 [M+Na]⁺ and HRMS *m/z* 1659.7831 [M+Na]⁺ (C₇₉H₁₁₂O₃₆). [α]_D²⁵ -14.3° (CHCl₃, *c* 0.35). ¹H NMR and ¹³C NMR data (CDCl₃): see Table 1.

3β-O-β-D-glucopyranosyl-(1-3)-α-L-arabinopyranosyl-20(S)-19α-hydroxyurs-12-en-28-oic acid 28-O-β-D-glucopyranosyl ester (4). Amorphous powder. FABMS (positive mode) *m/z*: 951 and HRMS *m/z* 951.4899 [M+Na]⁺ (C₄₇H₇₆O₁₈). [α]_D²⁵ -1.67° (MeOH, *c* 0.6). ¹H NMR and ¹³C NMR data (CD₃OD): see Table 1.

3β-O-β-D-glucuronopyranosyl-20(R)-19α-hydroxyurs-12-en-28-oic acid 28-O-β-D-glucopyranosyl ester (5). Amorphous powder. FABMS (positive mode) *m/z*: 833.6 [M+Na]⁺. ¹H NMR and ¹³C NMR data (CD₃OD): see Table 1.

RESULTS AND DISCUSSION

Solvent partition and chromatographic procedures allowed the isolation of five compounds from leaves of *I. chamaedryfolia* whose identification was accomplished through spectroscopic methods.

Compound **1** was identified as α-amyrin after comparison of its physical-chemical data with literature and by co-TLC with an authentic sample. Compound **2** was rapidly identified as ursolic acid by co-TLC analysis using authentic sample.

Along with the signals corresponding to acetyl groups, {¹H}-¹³C-NMR spectrum of **3a** showed 53 signals, which after comparison with its DEPT-¹³C-NMR spectrum, allow to recognize signals attributed to 7 methyl, 13 methylene and 25 methine groups, and 8 quaternary carbon atoms. The ¹³C-NMR of this compound showed characteristic signals due to one carboxyl group (δ 175.7), a double bond (δ 128.0; δ 137.0) and two oxygenated carbons (δ 72.9; δ 89.9) beyond the acetyl characteristic signals.¹² A signal at δ_H 2.77 (*s*, 1H, H-18) suggesting the presence of a 19-O-substituted urs-12-en skeleton, specifically a 19α-hydroxyursolic acid derivative.¹³ Chemical shifts of the C-18, C-22 and C-23 were found at upfield shift and indicate an unusual 20 *S* configuration to E ring of aglycone (Table 1).^{8,12} The observation of chemical shifts to C-28 (δ 175.7) and C-3 (δ 89.9) dislocated to upfield and downfield respectively, cleared bisdesmosidic features. Thus, the compound **3a** (and **3**) has as aglycone ilexgenin B (Figure 1).¹²

Its ¹H-NMR and ¹³C-NMR data indicated the presence of four sugar moieties. The sugar residue δ_H 5.55 (H-1''); δ_C 91.7 (C-1''') evidenced an ester linkage between the anomeric carbon and the C-28 of the aglycone.¹⁴ For the sugar chain, HH and HC correlation (COSY, HMQC and HMBC) (Figure 2) allowed to assign all carbon and proton

signals. HMBC correlation spectrum allowed indicated the interglycosidic linkages, between the terminal glucose (δ_H 4.73, H-1'; δ_C 99.7, C-1), attached at C-3''' of glucose II at C-28, as well as, glucose I and arabinose unit substituted at C-3. This spectrum showed a correlation between C-3' of arabinose and H-1'' of glucose I, C-3''' of glucose II and H-1'''' of glucose III (Figure 2). Besides H-1' of arabinose and C-3 of aglycone, H-1'' of glucose II and C-28 of aglycone. Thus **3a** was identified to be peracetylated form of 3β-O-β-D-glucopyranosyl-(1-3)-α-L-arabinopyranosyl-20(S)-19α-hydroxyurs-12-en-28-oic acid 28-O-β-D-glucopyranosyl-(1-3)-β-D-glucopyranosyl ester (Figure 1).




Figure 1. Compounds **3**, **4** and **5** isolated from leaves of *Ilex chamaedryfolia*




Figure 2. Main long range correlations observed in HMBC spectra for **3a** and **4**

Comparative analysis of the ¹³C-NMR-¹H and ¹³C-NMR-DEPT spectra of **4** was also used to identify the presence of signals attributed to 7 methyl, 12 methylene and 20 methine groups and 8 quaternary carbon atoms. The NMR data (¹³C; ¹H) of compound **4** were very similar to compound **3a** excepting the absence of the acetyl characteristic signals and those corresponding to one glucose. After detailed analysis of the spectroscopic data of **4** it was possible to conclude that this compound contained a single glucose residue in the ester chain at C-28, contrasting to **3a** and **3** that contained two glucose units. Thus, **4** was identified as 3β-O-β-D-glucopyranosyl-(1-3)-α-L-arabinopyranosyl-20(S)-19α-hydroxyurs-12-en-28-oic acid 28-O-β-D-glucopyranosyl ester (Figure 1). The HH and HC correlation (COSY, HMQC and HMBC) allowed to assign all carbon and proton signals and to identify all interglycosidic linkages (Figure 2). The 20R enantiomer of this compound has already been reported

Table 1. ¹H and ¹³C NMR spectral data for triterpenoids **3a**, **4** and **5** compared to literature

	3a , CDCl ₃		4 , CDCl ₃		5 , CD ₃ OD		Mo-1, CD ₃ OD	
	δ _c	δ _H	δ _c	δ _H	δ _c	δ _c	δ _H	
CH ₂ -1	38.6		38.5		39.9	39.90	1.64, 0.99	
CH ₂ -2	26.0		25.8		21.1	27.05	1.84, 1.73	
CH-3	89.9	3.10(m)	89.2	3.10 (m)	90.9	90.56	3.15 (dd, 11.5, 4.0)	
C-4	39.9		38.8		40.2	40.20		
CH-5	55.6		55.7		57.0	57.03	0.79	
CH ₂ -6	18.2		18.0		19.4	19.44	1.53	
CH ₂ -7	33.0		32.8		34.2	34.14	1.54, 1.33	
C-8	39.2		39.8		42.7	41.23		
CH-9	47.1		47.8		48.0	48.59	1.67	
C-10	36.6		38.8		37.8	37.84		
CH ₂ -11	23.8		23.3		24.6	24.73	1.97	
CH-12	128.0	5.35 (br s)	127.5	5.15 (br s)	129.8	129.66	5.33 (br s)	
C-13	137.0		137.8		139.5	139.50		
C-14	41.2		41.3		42.9	42.59		
CH ₂ -15	28.0		28.2		29.7	29.63	1.83, 1.02	
CH ₂ -16	25.8		25.7		26.5	26.51	2.16 (td), 1.63	
C-17	47.5		48.8		49.0	49.41		
CH-18	46.3	2.77 (s)	46.4	2.65 (s)	54.9	54.88	2.51 (s)	
C-19	72.9		72.9		72.9	73.63		
CH-20	41.4		41.8		41.3	42.87	1.36	
CH ₂ -21	23.6		23.6		27.2	27.19	1.71	
CH ₂ -22	30.6		30.8		32.2	38.24	1.77, 1.63	
CH ₃ -23	27.8	1.10 (s)	27.1	0.95 (s)	28.5	28.59	1.05 (s)	
CH ₃ -24	15.6	0.90 (s)	15.6	0.75 (s)	17.0	17.06	0.85 (s)	
CH ₃ -25	16.2	0.85 (s)	14.6	0.85 (s)	16.0	16.06	0.95 (s)	
CH ₃ -26	16.8	0.71 (s)	16.3	0.68 (s)	17.6	17.61	0.77 (s)	
CH ₃ -27	24.0	1.20 (s)	22.9	1.22 (s)	24.7	24.73	1.33 (s)	
C-28	175.7	-	177.1	-	178.5	178.46		
CH ₃ -29	29.9	1.30 (s)	28.9	1.06 (s)	28.9	27.14	1.20 (s)	
CH ₃ -30	15.2	1.00 (d, 6.0)	14.8	0.89 (d, 5.6)	16.6	16.63	0.93 (d, 6.8)	
sugars		3-O-Ara			3-O-GlcA			
CH-1'	104.2	4.30 (d, 6.9)	105.6	4.19 (d, 7.2)	106.7	107.01	4.29 (d, 7.0)	
CH-2'	72.9		69.7		73.9	72.04	3.71	
CH-3'	79.8		82.4		77.9	83.80	3.65	
CH-4'	72.9		68.1		75.5	69.46	4.03 (br s)	
CH-5'	-		-		78.3	-		
CH ₂ -5'	63.0		65.2		-	66.62	3.86 (dd), 3.55 (d)	
C-6'	-		-		180.8	-		
CH ₂ -6'	-		-		-	-		
		3-O-Glc						
CH-1''	100.6	4.72 (d, 7.1)	103.9	4.45 (d, 7.8)		105.30	4.56 (d, 7.3)	
CH-2''	70.0		73.9			75.25	3.31	
CH-3''	72.0		76.9			77.80	3.31	
CH-4''	68.0		69.8			71.13	3.37	
CH-5''	71.9		76.5			77.56	3.49	
CH ₂ -6''	61.5		61.0			62.34	3.82 (dd), 3.70	
		28-O-Glc						
CH-1'''	91.7	5.55 (d, 8.1)	94.4	5.25 (d, 8.2)	95.8	95.70	5.33 (d, 8.3)	
CH-2'''	70.2		72.5		73.9	73.78	3.33	
CH-3'''	73.9		77.3		78.6	78.45	3.35	
CH-4'''	67.9		70.7		71.1	71.05	3.37	
CH-5'''	72.4		76.3		78.3	78.20	3.42	
CH ₂ -6'''	62.9		61.0		62.4	62.39	3.82 (dd), 3.70	
CH-1''''	99.7	4.73 (d, 7.8)						
CH-2''''	69.9							
CH-3''''	72.0							
CH-4''''	68.0							
CH-5''''	71.9							
CH ₂ -6''''	61.5							

Mo-1 = 20(R) = randiasaponin III¹⁶ = ilekudinoside E¹⁵

in *Ilex kudincha*¹⁵ and *Randia formosa*.¹⁶ Thus, **4** is the epimer of ilekudinoside E and randiasaponin III.

In relation to **5**, through comparison of its ¹³C-NMR-{1H} and ¹³C-NMR-DEPT spectra, it was possible to identify the presence of 7 methyl, 10 methylene and 16 methine groups and 8 quaternary carbon atoms. ¹H-NMR joined to ¹³C-NMR data indicated the presence of two sugar residues. The later spectrum also evidenced the presence of one glucose residue linked to C-28 via an ester linkage (anomeric carbon at δ_c 95.8) and the presence of glucuronic acid moiety (δ_c 106.7) attached to C-3. Comparison of the NMR data of compound **5** to those of the literature allowed the identification of its aglycone as pomolic acid, one 19 α -hydroxyursolic acid derivative possessing a 20R-configuration.¹⁷ The β configuration for the glucopyranosyl and glucuronopyranosyl units and the α configuration for the arabinopyranosyl residues were inferred from their ¹³C-NMR data and *J* values.¹⁸ These data suggested that **5** is the 3 β -O- β -D-glucuronopyranosyl-20(R)-19 α -hydroxyurs-12-en-28-oic acid 28-O- β -D-glucopyranosyl ester (Figure 1) which was already isolated from *Ilex kudincha* by Nishimura *et al.*¹⁵ as ilekudinoside B.

To the best of our knowledge, the saponins **3** and **4** reported herein have not been yet described in the literature. Moreover, excepting few reports,^{8,19} the co-occurrence of triterpenes with configuration 20R and 20S is not a common feature to the *Ilex* species already studied. Indeed, it is possible to differentiate *I. chamaedryfolia* from other South-American *Ilex* considering their saponins.⁸

CONCLUSIONS

Several works have demonstrated that the saponins isolated from leaves of *I. paraguariensis* (maté) are different from those found in other South-American *Ilex* species.^{1,3-6,8} The aglycone pattern 19-hydroxyursolic and 29-hydroxyursolic acid found in saponins isolated from South-American *Ilex* species investigated up to now, allowed to differentiate all this species from *I. paraguariensis* leaves whose saponins are glycosides of ursolic or oleanolic acid. As suggested by our previous results,⁸ saponin content of South-American *Ilex* species can be used as a chemotaxonomic tool to the genera. This was confirmed herein considering that saponins so far isolated from *I. chamaedryfolia* are based on 19-hydroxyursane triterpenes which are different from those presented in the leaves of *I. paraguariensis*.

ACKNOWLEDGEMENTS

The authors are grateful to M. Sobral (Universidade Federal do Rio Grande do Sul) and S. A. L. Bordignon (Centro Universitário La

Salle, Canoas-RS) for collecting and identifying the plant material and to Unidade de Química de Proteínas e Espectrometria de Massas (Uniprote-MS) do Centro de Biotecnologia/UFRGS for HRMS. This work was supported by the Brazilian agencies: CNPq, CAPES and FAPERGS. The authors are also grateful to the Programa de Pós-Graduação em Ciências Farmacêuticas/UFRGS (Brazil).

REFERENCES

- Schenkel, E. P.; Gosmann, G.; Montanha, J. A.; Heizmann, B. M.; Athayde, M. L.; Taketa, A. T. C.; Pires, V. S.; Guillaume, D.; *Cien. Cult.* **1997**, *49*, 359.
- Giberti, G. C.; *Dominguezia* **1989**, *7*, 1.
- Pires, V. S.; Guillaume, D.; Gosmann, G.; Schenkel, E. P.; *J. Agric. Food Chem.* **1997**, *45*, 1027.
- Athayde, M. L.; Schenkel, E. P.; Gnoatto, S. B.; Gosmann, G.; Giberti, G. C.; Guillaume, D.; *Acta Farm. Bonaerense* **2001**, *20*, 13.
- Taketa, A. T. C.; Schlager, T. S.; Guillaume, D.; Gosmann, G.; Schenkel, E. P.; *Phytochemistry* **2000**, *53*, 901.
- Gosmann, G.; Schenkel, E. P.; Seligmann, O.; *J. Nat. Prod.* **1989**, *52*, 1367.
- Athayde M. L.; Schenkel, E. P.; Gosmann, G.; Guillaume, D.; *Acta Farm. Bonaerense* **1999**, *18*, 49.
- Pires, V. S.; Gosmann, G.; Guillaume, D.; Schenkel, E. P.; *Nat. Prod. Lett.* **2002**, *16*, 401; Gnoatto, S. C. B.; Bassani, V. L.; Coelho, G. C.; Schenkel, E. P.; *Quim. Nova* **2007**, *30*, 304.
- Edwin, G.; Reitz, P. R. In *Flora Ilustrada Catarinense. I Parte*; Reitz, P. R., ed.; Herbário Barbosa Rodrigues/CNPq: Itajaí, 1967, cap. Aquifoliáceas.
- Seo, S.; Tomita, Y.; Tori, K.; *Tetrahedron Lett.* **1975**, *1*, 7.
- The Merck Index: an encyclopedia of chemicals, drugs, and biologicals*; Budavari, S., ed.; Merck: Whitehouse Station, 1996, 12th ed., p. 1686-7.
- Hidaka, K.; Ito, M.; Matsuda, Y.; Kohda, H.; Yamasaki, K.; Yamahara, J.; Chisaka, T.; Kawakami, Y.; Sato, T.; Kagei, K.; *Chem. Pharm. Bull.* **1987**, *35*, 524.
- Wenjuan, Q.; Xiue, W.; Junjie, Z.; Fukuyama, Y.; Yamada, T.; Nakagawa, K.; *Phytochemistry* **1986**, *25*, 913.
- Massiot, G.; Lavaud, C.; Binst, V. G.; Miller, P. F. S.; Fales, H. M.; *J. Chem. Soc.: Perkin Trans. I* **1988**, 3071.
- Nishimura, K.; Miyase, T.; Nogushi, H.; *J. Nat. Prod.* **1999**, *62*, 1128.
- Sahpaz, S.; Gupta, M. P.; Hostettmann, K.; *Phytochemistry* **2000**, *54*, 77.
- Kakuno, T.; Yoshikawa, K.; Arihara, S.; *Phytochemistry* **1992**, *31*, 2809.
- Agrawal, P. K.; *Phytochemistry* **1992**, *31*, 3307.
- Ouyang, M. A.; Liu, Y. Q.; Wang, H. Q.; Yang, C. R.; *Phytochemistry* **1998**, *49*, 2483.