

DERIVADOS CITOTÓXICOS DE VITANOLIDOS ISOLADOS DAS FOLHAS DE *Acnistus arborescens*

Sandro Minguzzi*

Curso de Química, Universidade Estadual de Mato Grosso do Sul, Unidade de Naviraí, Rua Emílio Máscoli, 275, 79950-000 Naviraí - MS, Brasil

Lauro E. S. Barata

Instituto de Química, Universidade Estadual de Campinas, CP 6154, 13083-970 Campinas - SP, Brasil

Geoffrey A. Cordell

Natural Products Inc., 60203 Evanston - IL, USA




Figura 1S. Espectro de RMN ¹H de 2

*e-mail: sming@uems.br




Figura 2S. Espectro de RMN ^1H de 3




Figura 3S. Espectro de RMN ¹H de 2a




Figura 4S. Espectro de RMN de ^{13}C de 2a




Figura 5S. Espectro DEPT de 2a




Figura 6S. Espectro de RMN ^1H de **2b**




Figura 7S. Espectro COSY ^1H - ^1H de 2b




Figura 8S. Espectro de RMN de ^{13}C de **2b**




Figura 9S. Espectro DEPT de 2b




Figura 10S. Espectro de RMN ^1H de 2c




Figura 11S. Espectro de RMN ^1H de **3a**




Figura 12S. Espectro de RMN de ^{13}C de 3a




Figura 13S. Espectro de RMN ¹H de 3b