

SYNTHESIS OF GERANYLHYDROQUINONE DERIVATIVES WITH POTENCIAL CYTOTOXIC ACTIVITY

Evelyn Baeza, Karen Catalán, Hugo Peña-Cortés and Luis Espinoza*

Departamento de Química, Universidad Técnica Federico Santa María, Valparaíso, Chile

Joan Villena

Facultad de Medicina, Universidad de Valparaíso / Centro Regional de Estudios en Alimentos Saludables, Valparaíso, Chile.

Héctor Carrasco

Departamento de Ciencias Químicas, Universidad Andrés Bello, Campus Viña del Mar, Chile








Figure 1S. ¹H NMR (400 MHz, CDCl₃) of compounds 1

*e-mail: luis.espinozac@usm.cl

Figure 2S. ¹³C NMR (100 MHz, CDCl₃) of compounds 1Figure 3S. ¹H NMR (400 MHz, CDCl₃) of compounds 2

Figure 4S. ¹³C NMR (100 MHz, CDCl₃) of compounds 2Figure 5S. ¹H NMR (400 MHz, CDCl₃) of compounds 3






Figure 6S. ¹³C NMR (100 MHz, CDCl₃) of compounds 3



Figure 7S. ¹H NMR (400 MHz, CDCl₃) of compounds 4

Figure 8S. ¹³C NMR (100 MHz, CDCl₃) of compounds 4Figure 9S. ¹H NMR (400 MHz, CDCl₃) of compounds 5




Figure 10S. ^{13}C NMR (100 MHz, CDCl_3) of compounds 5