

SYNTHESIS AND BIOLOGICAL EVALUATION OF BIARYL ANALOGS OF ANTITUBULIN COMPOUNDS

Camila Santos Suniga Tozatti, Rejane Gonçalves Diniz Khodyuk, Adriano Olimpio da Silva, Edson dos Anjos dos Santos, Marcos Serrou do Amaral e Dênis Pires de Lima*

Centro de Ciências Exatas e Tecnologia, Universidade Federal de Mato Grosso do Sul, Av. Senador Filinto Müller, 1555, 79074-460 Campo Grande - MS, Brasil

Ernest Hamel

Screening Technologies Branch, Developmental Therapeutics Program, Division of Cancer Treatment and Diagnosis, National Cancer Institute at Frederick, National Institutes of Health, Frederick, MD 21702, USA




Figure 1S. EI-MS spectrum of 5a




Figure 2S. EI-MS spectrum of 3a




Figure 3S. EI-MS spectrum of 3f




Figure 4S. EI-MS spectrum of 3d




Figure 5S. EI-MS spectrum of 4b




Figure 6S. EI-MS spectrum of 3e




Figure 7S. EI-MS spectrum of 3h




Figure 8S. EI-MS spectrum of 4d




Figure 9S. EI-MS spectrum of 3g




Figure 10S. EI-MS spectrum of 4a




Figure 11S. EI-MS spectrum of 5c





Figure 12S. EI-MS spectrum of 3b




Figure 13S. EI-MS spectrum of 3c




Figure 14S. EI-MS spectrum of 5b




Figure 15S. EI-MS spectrum of 4c