

ISOLATION AND ANTITRICHOMONAL ACTIVITY OF THE CHEMICAL CONSTITUENTS OF THE LEAVES OF *Maytenus phyllanthoides* Benth. (Celastraceae)

Juan Alberto Moo-Puc^a, Zhelmy Martín-Quintal^a, Gumersindo Mirón-López^a, Rosa Esther Moo-Puc^b, Leovigildo Quijano^c and Gonzalo J. Mena-Rejón^{a,*}

^aLaboratorio de Química Farmacéutica. Facultad de Química. Universidad Autónoma de Yucatán. Calle. 41 No. 421 x 26 y 28, Col. Industrial. C.P. 97150, Mérida, Yuc., México

^bUnidad de Investigación Médica Yucatán, Unidad Médica de Alta Especialidad, Centro Médico Ignacio García Téllez, Instituto Mexicano del Seguro Social, 41 No. 439 x 32 y 34, Col. Industrial CP 97150, Mérida, Yuc., México

^cInstituto de Química, Universidad Nacional Autónoma de México, Circuito Interior, Ciudad Universitaria C. P. 04510, Coyoacán, México




Figure 1S. ¹H NMR spectrum of compound 1 (400 MHz, CD₃OD)




Figure 2S. ^{13}C NMR spectrum of compound I (100 MHz, CD_3OD)




Figure 3S. ^1H NMR spectrum of compound Ia (400 MHz, CDCl_3)




Figure 4S. ^{13}C NMR spectrum of compound **1a** (400 MHz, CDCl_3)




Figure 5S. ^1H NMR spectrum of compound **2** (400 MHz, CDCl_3)




Figure 6S. ¹H NMR spectrum of compound 3 (400 MHz, CD₃OD)




Figure 7S. ¹³C NMR spectrum of compound 3 (100 MHz, CD₃OD)




Figure 8S. ^1H NMR spectrum of compound 4 (400 MHz, CDCl_3)




Figure 9S. ^{13}C NMR spectrum of compound 4 (400 MHz, CDCl_3)